

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.



2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY IN: AL, AR, FL, GA, IN, KS KY, LA, MS, NC, OK, SC, TN & VA

THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL DIRECTIONS FOR USE WHICH ARE RECOMMENDED BY FMC CORPORATION UNDER FIFRA SECTION 2(ee).

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH THE LABELING.

CROP	PEST	RATE OF APPLICATION
Turf (Warm Season) Bermudagrass (<i>Cynodon dactylon</i>) & hybrids Centipedegrass (<i>Eremochloa ophiuroides</i>) St. Augustine grass (<i>Stenotaphrum secundatum</i>) Zoysiagrass (<i>Zoysia japonica</i>)	Poa Annua (Annual Bluegrass)	0.15 – 0.23 oz/1000 ft² 6.5 – 10 ounces/A 0.27 - 0.413 lbs ai/A

DIRECTIONS FOR USE: Applications may be applied to non-overseeded warm season turf in November/December, February –April for control of Poa Annua. Apply in 20 - 60 gallons of water /acre. Use higher rates for larger poa annua plant size and increased populations.

Application of this product can be made following the second mowing providing the turfgrass has developed into a uniform stand with a good root system. Turfgrass injury could result from application of this product on turfgrass that is not well established or has been weakened by stresses such as unfavorable weather conditions, disease, chemical or mechanical influences.

RESTRICTIONS: Do not apply to overseeded bermudagrass.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED BLINDSIDE HERBICIDE LABEL (EPA REG. NO. 279-3411) MUST BE FOLLOWED.

THIS RECOMMENDATION LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

